# JNIVERSITY GUELPH **Ontario Veterinary College**

**Health Sciences Centre** 

# **Companion Animal Surgery Service Discharge Summary**

Wendy Perkes wperkes@ovc.uoguelph.(q debbie@uoguelph.ca 605477



OVC HSC Clinicians: Dr. Brigitte A. Brisson (DVM, DVSc, Dipl. ACVS), Dr. Debbie Reynolds (BVSc)

Dupont, Amber 303 Riverside Drive Welland, ON L3C 5E6 H: (905) 732-2513 W: ( \_\_\_) \_-\_

Abby Canine Sheepdog, Old English 12/02/11 F Black & White Main West Animal Hospital Ph: 905-735-7877 Fax: 905-735-9800

#### Surgery Faculty and Staff Surgeon

**Referring Veterinarian:** 

Admitted On: 08/02/12

Released On: 12/02/12

In Clinic Summary

**Received By:** 

Diagnosis:

Sarah E. Boston DVM, DVSc, Dipl. ACVS

Brigitte A. Brisson DVM, DVSc, Dipl. ACVS

Thomas W.G. Gibson DVM, DVSc, Dipl. ACVS

Agatha Kisiel DVM, MSc

Noel M.M. Moens DVM, MSc, Dipl. ACVS, Dipl. ECVS

Ameet Singh DVM, DVSc, Dipl. ACVS

**Surgery Residents** 

Galina Hayes DWM, PhD, Dipl. ACVECC

Michelle Oblak DVM

Adam Ogilvie DVM

Debbie Reynolds BVSc

**Surgery Technician** 

Susanne Shearing RVT 519.824.4120 ext. 54449

Tests: Physical examination Upper gastrointestinal endoscopy Non selective angiography Thoracic radiographs

Vascular ring anomaly (persistent right ligamentum arteriosum)

## Treatments:

General anesthesia Intravenous fluid therapy Supportive care in the Intensive Care Unit Analgesia Antibiotics Prokinetic therapy Gastroprotectants

Surgery: Left exploratory thoracotomy Right exploratory thoracotomy and ligamentous arteriosum ligation and transection

## **Instructions to Owners:**

#### Exercise:

Please confine Abby to a small crate or pen for the next 2 to 3 weeks to allow her incisions to heal. Do not allow any running, jumping or rough play with her littermates during this time. The more active Abby is the more likely there are to be incisional complications including but not limited to swelling, infection and dehiscence (breakdown).

#### Diet:

Please continue to feed Abby gruel/slurry while standing on her hind legs with the food dish elevated on a platform. She should be maintained in an upright position (held upright in your arms) for 10 minutes after eating to allow gravity to assist in emptying the dilated esophagus. Our hope is that in 4